

# Pioneers of the Ice Age Models: A Brief History from Agassiz to Milankovitch

M. Efe Ateş<sup>1</sup>

<sup>1</sup>Department of Philosophy, MSKÜ, Mugla, 48000, Turkey

5 Correspondence to: M. Efe Ateş (mefeates@mu.edu.tr)

**Abstract.** It is now widely accepted that astronomical factors trigger the emergence of glacial and interglacial periods. However, nearly two centuries ago, the overall situation was not as apparent as it is today. In this article, I briefly discuss the astronomical model of ice ages put forward in the 19th and early 20th centuries. This period was indeed *anni mirabiles* for scientists to understand the ice age phenomenon. Agassiz, Adhémar and Croll laid the foundation stones for understanding the dynamics of ice ages. But it was Milankovitch who combined empirical geology with mathematical astronomy. Specifically, he identified the shortcomings of the preceding ice age models and modified his model accordingly. In what follows, I review earlier approaches to the ice age problem and show how they failed to meet their objectives. Next, I show how Milankovitch's model managed to capture all sufficient astronomical elements. The final sections focus on Milutin Milankovitch's successful approach, including his accomplishment of tackling the problem mathematically.

## 15 1 The Problem

There seems no strict quantitative definition of ice ages (see Davis, 2001; Barry and Yew Gan, 2011, p. 299). However, we will just say simply that ice ages are periods when severe temperature reduction occurs across the Earth. During such periods of time, the surface of the Earth is largely covered with ice sheets and mountain glaciers. So far, scientists provide evidence that the Earth, in its history, has experienced many glacial periods. However, only few of these have established dominance over the whole climate system by lasting more than millions of years. Fig. 1 shows these extensive ice ages in the past 2.4 billion years.

As seen from the chart, scientists have identified five significant ice age periods during the past 2.4 billion years. They are respectively called Huronian (2.4–2.1 billion years ago), Cryogenian (850–630 million years ago), Andean–Saharan (460–430 million years ago), Karoo (350–260 million years ago) and Quaternary (2.6 million years ago–present). Accordingly, we are living in the Quaternary period which began approximately 2.6 million years ago. Since we live in the Quaternary period, we are currently in the last major ice age period. So, the Earth should be in a notably colder climate compared to our current condition. At present, however, this is not the case. The climate is not extremely cold, nor are our continents largely covered by ice sheets. On what grounds, then, do scientists claim that we are in a major ice age period?

Geologists already have the answer to this question. It seems that the received view now is that the Earth currently is in an interglacial period that started approximately 15 thousand years ago (Lisiecki and Raymo, 2005). According to this view, our planet, sometime during an extensive ice age period, makes alterations toward warmer climatic conditions. After witnessing warmer climate conditions, it again goes into a deep freeze and this specific event recurs at intervals. Thus, in short, the Earth's climate is subject to certain quasi-periodic changes. On a geological scale these alterations are a normal progression, but for humankind a glacial-interglacial transition is a giant step.

At first sight, the intuitive way to evaluate the alterations between glacial and interglacial periods is to assume a probable mechanism behind it. The scientists did so as well. Whether periodic or not, the Earth has witnessed, and probably will continue to witness numerous glacial and interglacial periods. Thus, for scientists, the new task is to find out which factors are responsible for this. The common questions which they had dealt with were as follows: Why the Earth's climatic conditions change within major ice ages? What factors are involved in the separation of glacial and interglacial periods? Why do series of glacial and interglacial intervals vary in length? The remainder of this article will be concerned with these questions and the answers given to them, but before proceeding let me say few words about the occurrence of major ice ages.

## 2 Major Ice Ages

There exist many causes of major glaciation; including oceanic fluctuations, volcanic eruptions and surface albedo (see also Rohling *et al.*, 2012). Without the feedback mechanisms initiated by plate tectonics, however, long term oscillations of climate would be completely different. The main cause responsible to initiate an ice age period is plate tectonics. Alfred Wegener (1880–1930), the German geologist and meteorologist, has laid the groundwork for this idea. He spent his time primarily in Greenland and his field research was –to a certain extent– centered upon continental drifts that later led Harry Hess (1906–1969) to develop the revolutionary theory of plate tectonics. The theory of plate tectonics was not specifically proposed with the purpose of explaining the mechanism of major glacial periods. Nonetheless, it provided a useful framework to explain the dynamics of long term climate changes.

According to this theory, the separate continents today were once combined together. The well known supercontinent Pangaea existed approximately between 350–260 million years ago and different from today's positioning, its continental mass was mostly located in southern hemisphere of the Earth. The east part of South America and the west part of Africa were bound together where India was located to the southeast. After a considerable time, each of these landmasses broke apart and drifted away until reaching their current positions (Fig. 2).

This theory has much to say about geological phenomena. With its principles, for example, we can deduce hypotheses about other natural events, such as volcanic eruptions, earthquakes or processes of mountain formation. Eventually, all these phenomena are results of the movements of plates which are segments of the continents and the oceans.

Similarly, the beginnings and the ends of major glacial periods are also related with the movements of continents. Many relevant textbooks give a clue about this relationship by using the term 'continental glacier' because glaciers (whether they

are ice sheets or ice caps) can only form on ground. This information is crucial for the formation of glaciers, but it is also crucial for the explanation of the initial conditions of glacial periods. In order to understand this, it's sufficient to consider that the positions of continents have decisive influence over the global climate. To put it precisely, it happens as follows: When the continents are positioned mostly near the poles, a low amount of sunlight falls on a big portion of total continental area. This means that snow and ice accumulate over large areas. Hence, the albedo increases and a significant amount of sunlight returns back to space. As a result, global temperature decreases and ineluctably produces a glacial period. This is why, many researchers have asserted that "the centering of a land mass on the pole or grouping of land masses around a pole to be an essential condition for glaciation" (Hay 1996, p. 410).

This natural process can be exemplified by reviewing the structure of the supercontinent Pangaea. As I mentioned above, large portion of Pangaea continent was located close to the south pole of the Earth. This means that suitable conditions were provided for a glacial period to start. Today, based on paleoclimatological evidence, we know that the Earth experienced an extensive ice age about 350–260 million years ago, also the era when Pangaea existed.

At this point, it is especially important to note that the theory of plate tectonics has no room or explanation for the questions raised at the end of the previous section. It is because this theory accounts for the long-term changes of global climate and falls short of establishing short-term temperature variations in major ice ages. According to the theory of plate tectonics, plates move very slowly (1–1.5 centimeters per year on average), and due to this fact, continents which are parts of plates move slowly too. This basic mechanism of the drift is extremely gradual such that continents shift their positions over millions of years. So, any model deduced from the theory of plate tectonics appears to be capable of explaining long term climatic changes. However, it would unavoidably fail to provide an explanation of what factors may be responsible for our climate to warm up and regress back to a cold climate within major ice age periods. So, the factors that lead to major ice ages do not properly explain short-term climate changes. To find out the causes of the short-term climate variations within major ice ages, then, we must narrow down the time scale from millions of years to thousands of years.

### **3 An Astronomical Solution for a Geological Puzzle**

As described above, climatic changes within the ice age periods are not caused by movements or positioning of continents. For this reason, we need to seek the causes of these temperature changes, elsewhere, in the Earth system. Milutin Milankovitch (1879–1958), the Serbian mathematician, astronomer and engineer, thought so and came up with an idea to connect climatic changes with Earth's orbital variations. According to him, temperature changes during the major ice age periods depend on the different amounts of solar irradiance falling on the surface of the Earth. By all means, the amount of solar irradiance is driven not by the Earth's internal system but by an external force, namely the Sun. Therefore, a possible solution lies not only in the field of Geology, but also in the field of Astronomy.

The idea to understand the dynamics of ice ages from the astronomical perspective is not new, and hence it does not belong entirely to Milankovitch (see Andersen 1992; Hestmark 2018; Berger 2021, p. 1731). There are other details which make

Milankovitch an original figure in science. These details and his other contributions will be spelled out in more detail after. But first, I will outline briefly the pioneers of the astronomical/geological theory who influenced Milankovitch's approach to the ice ages one way or another.

#### 4 Pioneers of the Ice Age Theory

We know quite a lot about the glaciers and glaciation, such as how ice sheets move, how glaciers shrink and grow, or how glacial deposits are formed etc. The situation, however, was different over the past two centuries. In those days, few scientists were involved with the glaciation research. Moreover, many were not aware how crucial this natural phenomenon is in shaping the solid surface of the Earth.

The first sign of the ice ages was erratic boulders. These erratic boulders (large masses of rock) are found in places far from their bedrock source. In the early to mid-18th century, the received view was that these giant rocks are somewhere distant from their original places because they are transported and deposited by a great flood –as almost told in the Bible. Although he changed his mind later on, Charles Lyell (1797–1875), the British geologist, confidently defended such a view and accepted it as the most likely explanation for the emplacement and movement of boulders (Imbrie and Imbrie, 1979, pp. 21–22, Boylan 1998, pp. 156–158). While this view dominated the geological scene for over decades, it subsequently received criticisms from some researchers in the scientific community. The main objection urged against the flood theory was that such giant rocks are unlikely to be carried from low to high elevations via floodwater. So, there must be another mechanism at work to transport massive boulders from one place to another. In addition, there was one other observation that flood theorists had difficulty accounting for: scratched and grooved bedrock surfaces. These traces on the land surface were an indication of a serious problem because flood theorists were assuming a simple process of transportation of boulders from one place to another. However, in order to erode the underlying land surface such a way, boulders must strongly scrape against the ground that they move over. Therefore, floodwater alone was not a possible candidate to be the cause of scratched and grooved surfaces.

At the time when the level of the erratic boulder discussion manifested itself in a puzzling way, Louis Agassiz (1807–1873), the Swiss biologist and geologist, was working on fossil fishes. However, along with fossil research, his mind was busy with another issue: the possible causal link between erratic boulders and glaciation. According to Agassiz, the boulders which we find far from their bedrocks are in their current different places not because they are carried by the flowing water but because they are removed by an agent with much stronger force. His hypothesis was that the displacement of a boulder from its bedrock could most probably be produced by the effect of glaciers. Especially, for Agassiz, polished bedrocks were a tangible sign of an ice age that had taken place in the distant past. As he himself stated:

These surfaces are sometimes even, sometimes undulated, often traversed by furrows more or less deep and sinuous, but which never occur in the direction of the slope of the mountain. On the contrary, these furrows are oblique and longitudinal; in short, they have a direction which at once excludes the idea of a

125 current water having been the cause of the erosions... To any one who has seen the Alps, it is evident that  
it is the ice which has produced this polishing (Agassiz 1838, p. 177).

To tell the truth, this hypothesis was not entirely original (see Seylaz 2017; Davies 2017). The relationships between boulder  
displacement, polished bedrocks and glaciation were suggested by many travelers and scientists prior to Agassiz (see Berger  
2012, pp. 108–110). Given the history of science, however, this is not new news. It is a rare possibility that a scientific claim  
130 has not been expressed before in a similar way by any other scientist. However, there are different and even worse  
allegations about the Agassiz’s case. He has been accused of intellectual dishonesty for failing to disclose that his hypothesis  
was borrowed from Karl Schimper (1803–1867), a German botanist (see Evans, 1887). I will not delve into this story further  
because as the Latin proverb says *errare humanum est*. Although he allegedly did not cite or credit relevant previous work,  
we should emphasize or recognize his admirable attitudes and attributes. The initiative he took in the organization of the  
135 Swiss Society of Natural Sciences, for example, is worth appreciating.

In 1837, Agassiz gave a talk about his glacial theory to the Swiss Society of Natural Sciences at Neuchatel, although the  
members of the society were expecting to hear a talk about his new research results on fossil fishes. Instead, Agassiz took the  
opportunity and announced his new argument on the phenomenon on glaciation. Truly, from that moment onwards the  
dispute had begun. Agassiz was exposed not only to the criticisms of opposing camps, but also showed resistance to the  
140 career-oriented advices from the authorities of geology, including William Buckland (1784–1856) and Alexander von  
Humboldt (1769–1859). For example, von Humboldt once wrote in a letter:

I am afraid you work too much, and (shall I tell you frankly?) that you spread your intellect over too many  
subjects at once. I think that you should concentrate your moral and also your pecuniary strength upon this  
beautiful work on fossil fishes. In so doing you will render a greater service to positive geology, than by  
145 these general considerations... In accepting considerable sums from England, you have, so to speak,  
contracted obligations to be met only by completing a work which will be at once a monument to your own  
glory and a landmark in the history of science... No more ice, not much of echinoderms, plenty of fish...  
(Agassiz, 1886, pp. 267–272).

Despite such skepticism, only about 30 years later, the theory suggesting the link between erratic boulders and glaciation was  
150 widely accepted in the scientific community, and this constituted a milestone in the advancement of glaciation research.

After being established as a physical process that occurred once or many times in the past, glaciation became a subject of  
geological investigations. As Agassiz stated in a letter to his friend, the glacial periods are hence regarded “by geologists as a  
fixed fact” (Agassiz 1871/1949, p. 294). Now the leading issue was no longer about the existence of past glacial periods.  
Instead, it was about the initial conditions of glaciations.

155 The French astronomer and mathematician Joseph Adhémar (1797–1862), was the first prominent scientist who drew  
attention to the onsets of glaciation process. According to him, temperature changes (like glaciation–interglaciation) in both  
hemispheres occur due to the precession of the equinoxes. In his book *Révolutions de la Mer*, he argues as follows:

[B]ecause of the precession of the equinoxes, the sums of the number of hours of day and night in each hemisphere are unequal. This inequality produces a difference in the corresponding temperatures, and it is this difference that explains the origin of the ices at the two poles (Adhémar, 1842, p. 96).

As stated by Adhémar, the durations of day and night differ in each hemisphere (except two days of the year, the spring equinox on March 21 and the autumnal equinox on September 23). The axial tilt of the Earth is the key factor underlying this difference. Today, the north pole of the Earth points to Polaris, more commonly known as the North Star. However, this will not always be the case. For Adhémar, the axis of the Earth wobbles and changes orientation between pointing at Polaris and Vega with a cyclic period of 22.000 years. In other words, the North Star will be Vega about 11.000 years later. Because the orientation of the axis changes very gradually, the Earth's tilt remains almost unchanged over the period of a single year. Whichever hemisphere tilts towards the Sun gets more light and hence experiences warm seasons. For instance, the northern hemisphere tilts toward the Sun as the Earth begins to move away from the Sun (March 21–September 23). Conversely, the southern hemisphere tilts toward the Sun as the Earth begins to move closer to the Sun (September 23–March 21).

According to Adhémar, one other point needs to be considered as well i.e. Earth's elliptical path around the Sun. The information that the Earth orbits the Sun in an ellipse, not a circle had already been known by means of Keplerian laws since the seventeenth century. The first one of these laws states that the Sun is not located exactly at the center of the ellipse, but rather located at one of the two foci. When the Earth gets closer to the Sun, the speed of our planet increases due to gravitational motion. The orbital speed of the Earth affects the number of days in each season. For example, the cold seasons (winter–autumn) of the northern hemisphere are short in number of days. On the other hand, when the Earth moves away from the Sun, the gravitational pull becomes weaker and our planet travels on its orbit at a relatively slow speed. As the Earth's orbital motion slows down, the northern hemisphere experiences warm seasons (summer–spring). Thus, the warm seasons of the northern hemisphere are more in numbers of days. Therefore, in the northern hemisphere, the number of days in the warm seasons is more than the number of days in the cold seasons.

According to Adhémar, the difference in the number of days between the two seasons exactly corresponds to 168 hours, i.e. 7 days. The situation is reversed, however, for the southern hemisphere. “The southern [hemisphere] will... lose, in a year, more heat than it receives, since the total duration of its nights exceeds that of the days by 168 hours” (Adhémar 1842, p. 27). Following this calculation, Adhémar concluded that the northern hemisphere enjoys longer summers, while the southern hemisphere experiences longer winters. Because the southern hemisphere experiences longer winters, it is actually in the process of glaciation.

At the time Adhémar's view were widely criticized. Many researchers, especially his contemporaries, “did not want to believe that the southern hemisphere is on average much colder than the northern hemisphere” (Bard 2004, p. 632). Despite these criticisms however, Adhémar still deserves to be remembered as a leading figure in the history of science, for laying the foundations of the astronomical theory of ice ages.

Another remarkable attempt to understand the onsets of glacial periods was made by the Scottish scientist and geologist James Croll (1821–1890). Croll was aware of Adhémar's work, and was also sympathetic to some parts of it. He, for

example, believed that long and cold winters would lead to glaciation in the corresponding hemisphere, as Adhémar suggested. Nevertheless, he was not inclined to ascribe particular importance to the precession of the equinoxes as a cause of glaciation. As we can recall, Adhémar was correct in stating that the northern hemisphere has long warm seasons, while the southern hemisphere has long cold seasons. In this fashion, he concluded that the northern hemisphere was experiencing an interglaciation, since it receives a great deal of insolation annually. On the other hand, the southern hemisphere was experiencing a glaciation due to opposite circumstances. However, Croll thought that this conclusion is wrong because if a particular hemisphere, for example, receives less sunlight in cold seasons, it would in turn receive more sunlight in warm seasons. Thus, the amount of heat loss in the winter season will be compensated by the following summer season in both hemispheres. As Croll puts,

Whatever extra heat the southern hemisphere may at present receive from the sun daily during its summer months owing to greater proximity to the sun, is exactly compensated by a corresponding loss arising from the shortness of the season; and, on the other hand, whatever daily deficiency of heat we in the northern hemisphere may at present have during our summer half-year, in consequence of the earth's distance from the sun, is also exactly compensated by a corresponding length of season (Croll, 1875, p. 82).

According to Croll, the compensation of heat loss or heat gain would almost nullify the effect of the axial precession. In this regard, no significant climatic impact would occur in each hemisphere. Thus, he reasoned that glacial periods must occur in periods when the heat compensation fails between cold and warm seasons. Axial precession alone might be a necessary factor, but apparently it was not sufficient. Therefore, there must be another factor at play to prevent the heat compensation mechanism from functioning. Croll identified this factor as high eccentricity. This also was the very idea which separates Croll further from Adhémar. As stated above, Adhémar knew that the Earth revolves around the Sun in an elliptical orbit. Nevertheless, he neglected the fact that this orbital shape might slightly change in time. In other words, he took eccentricity fixed at some value between 0 and 1 (circle and parabola, respectively). According to Croll, on the other hand, the modulation of axial precession by eccentricity should be considered because the mutual heat compensation between seasons could fail only in such condition. In this sense, he stated that we should take the combined influence of these two causes into account,

There are two causes affecting the position of the earth in relation to the sun, which must, to a very large extent, influence the earth's climate; viz., the precession of the equinoxes and the change in the eccentricity of the earth's orbit. If we duly examine the combined influence of these two causes, we shall find that the northern and southern portions of the globe are subject to an excessively slow secular change of climate, consisting in a slow periodic change of alternate warmer and colder cycles (Croll 1864, p. 129).

By means of two orbital factors, Croll explained the conditions in which the seasonal temperature difference could not be compensated for a given hemisphere. The hemisphere which tilts away from the Sun at times when the Earth is located at the far end of the elliptical orbit with a high eccentricity, it would likely to be in a process of glaciation. Croll claims that these two factors are responsible for unstable conditions of the seasons in a given hemisphere; nonetheless, they are not the *direct*

causes of glaciation. Put it more precisely, it is the Earth's physical agents that produce glaciation processes, not factors such as orbital forcing. These physical agents comprise many phenomena from the Earth's surface to Stratosphere, including clouds, fogs, wind-driven ocean currents etc. Croll, for example, described the effects of certain meteorological factors on glaciation as follows:

230 Under a cloudless sky, the direct rays of the summer-sun would ... be more than sufficient to remove the winter's accumulation of ice and snow. But if from thick fogs or an overcast sky the direct rays of the sun were prevented from penetrating to the earth, the heat of summer would not in such a case be sufficient to remove the snow and ice; and the formation of glaciers would be the inevitable result (Croll 1864, p. 133).

Among the phenomena that Croll identified as physical agents, the most noticeable one is the albedo effect. It is also  
235 noteworthy that no scientist until his time drew attention to this phenomenon (see Bol'shakov, Kapitsa and Rees 2012; Thompson 2021). The albedo effect is a mechanism that triggers the glaciation by reflecting the solar radiation back to space. As glaciers spread across landmasses, they cause the Earth to absorb less heat. In due course, the average temperature of our planet starts decreasing. This, in turn, creates a positive feedback loop which promotes the further spread of glaciers till they reach their maximum level. Here, one may easily identify the circular process. However, in addition to being  
240 circular, this process is also self-reinforcing. The effect amplifies the further effect of its initial cause, or to sum up in Croll's own words, "cause and effect mutually react so as to strengthen each other" (Croll, 1875, p. 75).

As it seems, Croll attributes the primary cause of glaciation to physical agents (e.g. albedo, ocean circulations). Put differently, the Earth's orbital motions only set the stage and conditions for physical agents. Nevertheless, it is worth mentioning that for Croll, there is no fundamental distinction between causes and conditions. This view was expressed in his  
245 highly philosophical work, published a few weeks before he died:

The distinction between cause and conditions is to a great extent arbitrary. There is no real or essential distinction between the two ... [A]ll the conditions are co-operating causes; and the selected one, which we term the cause, is effective only in co-operation with the others (Croll 1890, p. 68).

I believe this quote is key to understanding Croll's general view on the relation between glaciations and orbital motions. In  
250 this sense, physical agents (like albedo) are directly responsible for causing great climatic changes; however, orbital motions of the Earth are responsible for determining the appropriate initial conditions. As he states: "an increase of eccentricity could have no direct tendency to lower the temperature and cover [a] country with ice, [but] it might bring into operation physical agents which would produce this effect" (Croll, 1875, p. 13).

Agassiz, Adhémar and Croll are three significant figures that influenced the course of the glaciation research. However, they  
255 are not the only ones who deserve credit. There are many other scientists who contributed substantially to the field of ice age studies, by mainly observing nature over the years and making hypothesis about the past, present and future of the environmental conditions. Particularly, Imbrie and Imbrie (1979), Bard (2004), Berger (2012) and Paillard (2015), in their seminal and detailed works, traced the history of great glaciologists, their naturalist approaches and the breakthroughs in



glaciation research. In the following section, however, I focus on Milankovitch whose contribution had a significant impact  
260 on the science of glaciation among all those researchers.

## 5 The Astronomical Model of Ice Ages

Like many scientists, Milankovitch stood on the shoulders of giants. The contribution he made to science, particularly to  
climatology was only possible with the past contributions of great scientists. Indeed, Milankovitch often mentions the names  
of his predecessors in his book *Kanon der Erdbestrahlung und seine Anwendung auf das Eiszeitenproblem*, not only to  
265 reference their past research work, but also to give credit to their scientific legacy.

Nevertheless, as much as the past efforts should be recognized, we should also emphasize Milankovitch's unique approach  
to the problems. His approach is different from most of his predecessors because it involves some sort of unifying  
framework. Until the era of Milankovitch, the mainstream methodology on particular issues in geology was dominantly  
descriptive. However, for Milankovitch exact methodology (or in other words, mathematical approach) should be integrated  
270 into this descriptive stance. For him, only if this condition is met, could we hope to discover the geological traits of the Earth  
and other planets.

The time when Milankovitch encountered the problem concerning the glacial and interglacial periods, he was already  
equipped with this line of thought. According to him, those who were dealing with the issue had the necessary information to  
proceed, but they were not able to achieve it. Of course, there have been reasons for that. Given Milankovitch's  
275 interpretation, some of the scientists had little idea what to do with empirical evidence and how to relate them with relevant  
theory; while the others were unable to use their theoretical knowledge to construct the model of the investigated  
phenomenon (see Imbrie and Imbrie, 1979, pp. 97–99; Petrovic, 2012). In all this, Milankovitch was defending a unified  
view that suggests a combination of both approaches. Although, scientists like Croll made similar attempts to unite these  
approaches, they inevitably failed due to lack of empirical evidence or their insufficient mathematical training (see  
280 Milankovitch, 1941, p. 376).

As stated above, the way Milankovitch approached the problem was quite original. He argued that crucial climatic changes,  
such as series of glacial and interglacial intervals, are causally connected with the distribution of solar radiation that reaches  
the Earth's surface. Thus, any change in the amount of insolation triggers a change in the global climate. So, in order to  
understand much about the periods of glacial and interglacial climates, we must determine *all* the factors that vary the  
285 amounts of insolation.

According to Milankovitch, the distribution of incoming solar radiation happens in accordance with the Earth's orbital  
variations. These orbital variations cause little or no change in the amount of total insolation that reaches the Earth's surface,  
but they do play an essential role in seasonal changes. For example, they change the seasonal duration (long or short  
winters/summers) or the degree of difference between the seasons (hotter, milder or cooler winters/summers). Moreover,

290 these orbital variations comprise three astronomical cycles: orbital eccentricity, axial tilt and precession. Most commonly they are called ‘Milankovitch Cycles’ in deference to the Serbian scientist.

## 6 Milankovitch Cycles

I have briefly discussed two orbital elements (axial precession and orbital eccentricity) within the context of Adhémar’s and Croll’s approach. Nevertheless, there is a third orbital element called obliquity in addition to these two. In this section, I will  
295 briefly explain how each of these three elements operates in a cyclic fashion. The orbital movements of the Earth are referenced multiple times throughout the *Kanon*. Nevertheless, it is in the fourteenth chapter that Milankovitch represents all three cycles in the form of a compact diagram (Fig.3).

In the diagram “S represents the centre of the [Sun], and the ellipse PIAIIP [represents] the annual terrestrial orbit round the Sun” (Milankovitch 1941, p.246). The line SV drawn through S is normal to the orbital plane and “the angle VSN represents  
300 the inclination of the rotational axis of the Earth or the obliquity  $\varepsilon$  of the ecliptic” (Milankovitch 1941, p. 246). P and A respectively denotes the perihelion and the aphelion. Finally “the angle IISP, which may be called  $\gamma$ , represents the longitude of the perihelion to the vernal point” (Milankovitch 1941, p. 246). Fig. 3, in its original form, perfectly illustrates the dynamics of the Earth’s orbital motions but to provide further understanding, let me deconstruct the diagram into its three components and focus on each cycle separate from the others. (Fig. 4)

305 The first component of orbital variations is called orbital eccentricity. It can simply be described as the Earth’s orbital path around the Sun. The gravitational tug of other large planets influences Earth’s orbit. This leads to a change in orbital shape and hence produces a cycle lasting about 90000–100000 years. Actually, our planet appears to orbit around the Sun elliptically. However, the shape of this orbit is not fixed. Sometimes it becomes more elliptical and sometimes it becomes more circular. The orbit becomes more elliptical when the difference between the furthest and closest distance of the Earth  
310 from the Sun increases. Differently, when the difference between the furthest and closest distance of the Earth from the Sun reduces, the orbit becomes more circular. The place where the Earth is nearest to the Sun is called perihelion, which occurs around January 3. On or around July 4, however, the Earth is at its greatest distance from the Sun, which is called aphelion (Fig. 4a).

Orbital eccentricity takes a value obtained by a simple formula using the variables aphelion (a) and perihelion (p) as follows:  
315  $e = (a - p)/(a + p)$ . If the obtained value for orbital eccentricity is equal to zero ( $e=0$ ), the shape of the orbit is a perfect circle. If it is greater than zero ( $e > 0$ ), then the shape of the orbit is an ellipse. When the current estimated values are inserted into the above equation, we get  $e = 0.016$ .

Orbital eccentricity has an influence on climate change. However, it is the least effective factor on glaciation, among other variations. The reason is that the eccentricity variations have small impact on total annual insolation, namely a difference of  
320 0.03% (see Maslin and Ridgwell, 2005, p. 21). Nevertheless, orbital eccentricity of the Earth may lead to significant temperature contrast between seasons. For example, if the orbital path of the Earth around the Sun was circular, there would

be no annual insolation difference between summer and winter for each hemisphere. But, due to gravitational effects of other celestial objects, especially Jupiter's, this cannot happen. As stated, today, our planet's path is elliptical in shape, approximately with a value of 0.016. Therefore, the difference of seasonal insolation between summer and winter reaches to about 6%. So, this shows that the amount of solar radiation reaching to the Earth's surface is greater at perihelion, for northern hemisphere. If the Earth's orbital eccentricity is at its maximum value (0.07), namely the most elliptical shape, difference of seasonal insolation between summer and winter would reach to about 30%. As a result, seasons would be at their extremes in terms of temperature (e.g. very hot summers) for one hemisphere and would be moderate (e.g. milder summers) for the other.

The second component of orbital variations is called axial tilt or obliquity. Our planet's rotational axis is tilted relative to its orbital plane. The angle of the tilt is determined by drawing a line perpendicular to the Earth's orbital plane. The angle of the tilt changes between  $21.5^{\circ}$  to  $24.5^{\circ}$  on a 41000 year cycle (Fig. 4b)

Today, the obliquity of Earth is measured as  $23.5^{\circ}$ . This tilt can be measured easily at solstices and equinoxes. In order to do that, it is sufficient to take the inverse tangent of the value which is found by dividing an object's height by its shadow length, at that particular time. When the angle of tilt is about to increase its maximum value  $24.5^{\circ}$ , the temperature contrast grow sharper between the two seasons. In such a case, winters become colder and summers become warmer. Contrarily, when the angle of tilt decreases, the characteristics of seasons come closer. In this case, milder winters follow cooler summers.

The third and last component of orbital variations is called precession of the equinoxes. While orbiting the Sun, our planet also wobbles on its axis, like a spinning top. This wobbling of the Earth on its axis periodically repeats itself every 26.000 years. Today, the polar axis of the Earth points to Polaris, also known as the North Star. In fact, this won't last forever. Due to wobbling motion, the Earth's axis will gradually change and it will point to Vega. More specifically, around 13.000 years from now, Vega will be the North Star and 13.000 years after that Polaris will once again be the North Star (Fig. 4c). Thus, in effect, it takes 26 000 years for the axis to return to the same spot on the circle.

The main effect produced by precession is an alternation of the seasons for each hemisphere. For example, today, the northern hemisphere is in winter, the southern hemisphere is in summer at perihelion. An opposite situation will be observed by the middle of the cyclic period. To put it differently, when the Earth's rotational axis points to Vega, the northern hemisphere would be in summer and the southern hemisphere would be in winter at perihelion. In such a situation, the summers become warmer and the winters become colder. In other words, seasonal contrast in temperature becomes intensified.

While the notion of orbital cyclicity was not new at all for many astronomers, there was no available calculation of past orbital variations until the work of Urbain Le Verrier (1811–1877), the French astronomer and mathematician. By applying Newtonian laws to the masses of planets, he calculated the past changes of orbital motions of the Earth. In this way, he created a sort of data table that displays the past variations of Earth's orbit in the last 100000 years. After decades, American astronomer John Nelson Stockwell (1832–1920) calculated the changes in obliquity and eccentricity of eight planets in the

solar system. Furthermore, in 1904, German mathematician Ludwig Pilgrim (1844–1927) extended the calculations further and provided data on orbital changes of the Earth for the last million years.

Having these data was a big step forward for Milankovitch because they were used in revealing the relationship between insolation amount and global climate change of the Earth. However, Milankovitch was extremely careful and meticulous in his endeavors. He knew that the changes in solar irradiation are causally connected with the three orbital variations. He was also perfectly aware that these variations could be determined by precise calculation of all the planets' masses and of their motions. So, had they not inserted the actual masses of the planets while calculating the past orbital variations, the amount of insolation received by the Earth could not have been calculated correctly for remote past (see Milankovitch, 1941, p. 252–253).

Milankovitch firstly made use of the calculations done by Pilgrim. The reason was two-fold. First, Pilgrim's calculations covered the past 1 million years of the Earth's orbital variations. Using a calculation with such a wide time interval would certainly make things easier in understanding the past insolation amounts of the Earth. Second, Pilgrim's calculations were done including the times when Earth's longitude of perihelion was equal to  $90^\circ$  and  $270^\circ$ . The importance of this is that, according to Milankovitch, great climate changes tend to take place at times when the terrestrial perihelion attains the values of  $90^\circ$  and  $270^\circ$ .

Pilgrim was a good mathematician, or so Milankovitch had heard (see Milankovitch, 1941, p. 253), but he made use of Stockwell's integral formulae which contained many printer's errors as well as calculation mistakes, as first pointed out by German mathematician and astronomer Paul Harzer (1857–1932). In such a condition, all that remained was the calculations of Le Verrier's. Compared with the works of Pilgrim and Stockwell, Le Verrier's calculations were better, but unfortunately they were also somewhat inaccurate. Milankovitch, in a sense, was stuck for an answer and did not know what kind of strategy is best for his purposes. Eventually, critical support came from his colleague Vojislav Miskovitch (1892–1976), the Serbian astronomer. Using Le Verrier's integrals, Miskovitch did all the necessary re-computation by correcting the mass values of the planets (see, Janc *et al.*). This was thought to be a highly effective way because it would allow them to make a reasonable comparison, particularly between Le Verrier's recomputed results and Stockwell–Pilgrim's results. If the results indicate a slight difference, then the Earth's past insolation values could reliably be derived from these –otherwise things could get really tricky. Luckily, this strategy took hold in a significant manner. “The comparison of the insolation values that Milankovitch calculated from these solutions show[ed] a good agreement” (Berger 2021, p. 1728). After settling these preliminary points, the remaining task was essentially to combine the orbital variations of our planet with a mathematical model.

Milankovitch was well aware of the past astronomical theories and their shortcomings in explaining the phenomenon of glaciation. For example, Adhémar correctly emphasized the 26.000 years (however, Adhémar actually thought it was 22.000 years) cycle of precession as one of the most responsible factor for glaciation but considered the orbital eccentricity as a constant. Hence, this incorrect thought, no doubt, led him to conclude that ice ages could occur only in one hemisphere. In other words, his theory suggested that one hemisphere would experience ice age conditions while the other would be ice–

390 free during a period of 13.000 years (again, for Adhémar it was 11.000 years). Croll did not make the same mistake and paid much attention to the Earth's orbital eccentricity. Furthermore, he was the first who noticed the idea of albedo, or the reflectiveness of the Earth's surface. Although the effect of albedo is decisive to some extent, unfortunately that idea led him to think that colder winters would influence the expansion of glaciers. Thus, apart from neglecting the effect of obliquity, he insisted that ice ages are driven by very cold winters at aphelion.

395 According to Milankovitch, both earlier theories were incomplete because they do not contain all necessary parameters. In other words, either one or two orbital parameters are used in these early theories to give an explanation for glaciation. However, all parameters have an effect on the glaciation, one way or another and they should all together be taken into account. In order to be a significant global change in the climate, all the cyclic variations of the orbital motions must be in a state of superposition (see Milankovitch, 1941, p. 271). Because all three cycles operate independently "sometimes their  
400 influence on the amount of heat received by certain parts of the Earth nullifies each other" and "sometimes the changes increase or decrease the quantity of the heat" (Grubic, 2006, p. 199). The complex nature of orbital variations makes it difficult to express them mathematically, and it was precisely this difficulty that plagued Milankovitch's predecessors. According to him, both early approaches were mathematically incapable of linking the effects of orbital variations on insolation. He states this explicitly in his book *Kanon* as follows:

405 All these theories have ... the same shortcoming: None of them have correctly grasped the variability of all the astronomical elements which affect the irradiation of the Earth ... [and] besides, none of these theories was able to tackle mathematically the decisive influence of variations in the obliquity on the irradiation of the Earth (Milankovitch, 1941, p. 376).

And then continues as follows:

410 I was able to show that the astronomical problem of ice ages is far more complicated than had been assumed before, and that in order to arrive at a correct solution the whole problem had to be approached fundamentally and put on a broad basis (Milankovitch, 1941, p. 376).

Milankovitch's fundamental approach was to base his work on a mathematical model. Before constructing the model, he set some preliminary ground for a concept, namely the canonic unit (see Grubic, 2006, p. 199). Early on his research,  
415 Milankovitch noticed something interesting and important about climatological works. Until his time, no one had attempted to propose a mathematical theory of climate. One of the main reasons for this was that the majority of climatologists gave prime importance to empirical studies. The confidence in their instruments (with which they could make accurate measurements) was far greater than their confidence in a theory (which may turn out to be inaccurate). Another reason was related to a natural phenomenon, namely the Sun's rays. Nearly everybody knew that the Sun is the Earth's largest source of

420 radiant energy, and that also its rays pass through the atmosphere. What was unknown was whether the strength of these rays could be expressed as a constant.

American astrophysicist Samuel P. Langley (1834–1906) was the first scientist who initiated the research on solar constant at the Astrophysical Observatory of Smithsonian Institution. Langley's aim was to measure the mean solar radiation amount with the bolometer, an instrument he had invented. In this way, the Sun's rays could be expressed in calorie per square  
425 centimeter per minute. In 1884, Langley calculated the solar constant radiation corresponding to 3.1 grams–calories per cm<sup>2</sup> per minute (see also Langley, 1903). Almost thirty years after, however, in the yearbook of the same institute, scientists announced that they determined the new value of solar constant corresponding to 1.946 gram–calories per cm<sup>2</sup> per minute. Although Langley was not able to establish solar constant precisely, this unit is named *langley* (Ly) to honor his legacy.

Milankovitch was aware of these studies at the Smithsonian Institution (see Milankovitch, 1941, p. 306). Thus, he knew that  
430 the mean value of solar constant is found to be 1,946 grams–calories per cm<sup>2</sup> per minute. It was clear, for Milankovitch, that decades were spent to obtain the solar constant value; nevertheless, it could not be determined precisely. So, “this value [1,946 grams–calories per cm<sup>2</sup> per minute] is not yet to be considered as final” either (Milankovitch, 1941, p. 213). It seemed reasonable for Milankovitch not to use a value whose precision has not yet been established. For this reason, he expressed the mean value of intensity of the solar radiation in canonic units, and considered canonic unit ( $J_0$ ) as the solar constant. He  
435 took canonic unit “as the unit of radiation, and a hundred thousandth part of the year as the time unit” (Milankovitch, 1941, p. 266). To put it in symbols,  $J_0=1$ ;  $T=100.000$ .

It may seem perhaps arbitrary to express average solar radiation in canonical units, but actually it is not. According to Milankovitch, using the canonical unit would be advantageous to compute the course of insolation, i.e. winter and summer half–year insolation ( $W_w$ ,  $W_s$ ) and thus total year insolation ( $W_T$ ). In his words, “in computing the secular variations of  $W_w$ ,  
440  $W_s$ ,  $W_T$  and the secular march of terrestrial irradiation, it will be particularly advantageous to express insolation in question in canonic units” (Milankovitch, 1941, p. 266). To illustrate this computational advantage, it is sufficient to refer to two separate tables in *Kanon*, particularly Table VII and Table XI. In both tables, the insolation values of  $W_w$ ,  $W_s$ , and  $W_T$  are expressed in canonical units for the latitudes from 0° to 90°. In these tables, it is not specified to which hemisphere a particular latitude belongs to because, according to Milankovitch, for example, 65° N and 65°S receive the same amount of  
445 irradiation in their northern and southern summer half–years, respectively.

$W_w$ ,  $W_s$ , and  $W_T$  of the particular latitude differ between the two tables because they are produced with the same equations but different coefficients. Table VII, for instance, show the total year radiation quantities of 65° by using the heat unit  $J_0 = 2$  (notice that, this value is very close to 1.946) and the time unit 365, 2422 days, i.e. 525.942 minutes. Thus, the coefficient  $TJ_0$  in Milankovitch's initial insolation model is 1.051898 ( $2 \times 525.942$ ). On the other hand, Table XI indicates  $W_T$  of 65° in  
450 terms of canonic units, so  $TJ_0$  in that model is 100.000 ( $1 \times 100.000$ ). When these calculations are made, by taking into account the relevant orbital parameters, the insolation amount for 65° in Table VII and Table XI is respectively as follows:

Table VII 65° ( $W_s=143.000$ ,  $W_w=22.180$ ,  $W_T=165.180$ )

Table XI 65° ( $W_s=13.594$ ,  $W_w=2.109$ ,  $W_T=15.703$ )

All the values shown in Table VII can also be obtained from Table XI. To do this, it is necessary to divide the coefficient  
455 value in the equation used for table VII by the coefficient value in the equation used for Table XI ( $=1.051898/100.000$ ).  
When the obtained number 10.519 is multiplied by  $W_T$  of  $65^\circ$  in Table XI, we get approximately  $W_T$  of  $65^\circ$  in Table VII, i.e.  
165.180 ( $10.519 \times 15.703 = 165.179$ ).

All these calculations actually show one thing: Milankovitch's simplifications are valid. The time unit is not the actual  
number of minutes in a year (525,942 minutes). Rather, a year is divided into 100,000 equal parts. The value of heat unit is  
460 not 1.946 grams–calories, but is accepted as  $J_0=1$ . Despite all these simplifications, however, the insolation model using the  
canonical unit is in great agreement with the model that contains actual units. Moreover, the simplified model reduces the  
calculation labor as well.

After having settled all these matters to his satisfaction, Milankovitch presented his mathematical model for insolation. The  
essential elements for the model are as follows: summer and winter half–years for a certain year, changes of irradiation at a  
465 certain latitude for summer and winter, the change of the inclination of the ecliptic, eccentricity of the ecliptic in the given  
year, longitude of the perihelion relative to equinox and the coefficient for individual latitudes.

So, the model comprises all the necessary components including three orbital parameters, the changes of irradiation amount  
for specific latitudes and the coefficient that expresses the insolation amount in canonic units. With all these, the  
mathematical model would allow us to calculate the secular march of insolation for both the northern and southern  
470 hemispheres.

To understand how much solar radiation that the Earth received in the past thousands of years, we need to know what the  
current insolation value is. If the model can accurately provide the current insolation values, by relying on the actual values  
of the orbital parameters, then, its results can be extended to represent the past and as well as the future climatic conditions  
of the Earth. Thus, the retrodictive and predictive inferences about the Earth's climatic patterns would be valid if and only if  
475 they are deduced from true statements, i.e. from verifiable or factual statements. In light of these considerations,  
Milankovitch believed that his mathematical model needs to be verified or tested through empirical evidence. For this  
reason, he decided to calculate the mean annual temperature of the lower layer atmosphere for various latitudes. If the  
calculated results significantly agree with the empirically measured data, then the model would pass a test of verification.

Weather stations, as they do today, were regularly measuring the mean annual temperatures at various latitudes.  
480 Observations of weather conditions were recorded and transformed into data outputs. Milankovitch planned to compare the  
results of his model with this available data at his time. Focusing particularly on the northern hemisphere, he performed  
certain calculations. By means of the model, he assessed trends in average temperatures for northern latitudes. The average  
annual temperature values of the northern latitudes between  $40^\circ$ – $50^\circ$  were almost the same as the average temperatures  
recorded by the weather stations. However, there were significant temperature deviations at latitudes from  $40^\circ$  N to the  
485 equator and from  $50^\circ$  N to the north. Milankovitch, for example, found the average temperature of  $10^\circ$  N to be higher than the  
value determined by the weather station. The opposite held also for the upper latitudes of the northern hemisphere. This  
situation in fact weakens Milankovitch's model, rather than verifying it.

Nevertheless, Milankovitch anticipated the factors that led to the difference or failure: air and ocean currents. He reasoned that air and ocean currents carry canonic units from one place to another. Therefore, canonical units do not actually disappear; so they are (or they should be) stored somewhere in the Earth's atmosphere. Accordingly, Milankovitch revised his approach to the topic, by calculating the mean temperature of the lower layer atmosphere for all the Earth's latitudes. The new results are compared with those of weather stations. The mean value difference between Milankovitch's model versus meteorological measurements was only 0.1° C. To put it clearer, this very slight difference indicated that Milankovitch's model is also verified or at least highly supported by empirical evidence.

With the Milankovitch model of insolation, it became possible to calculate how much solar radiation is received for any latitude in the past, present and future. Nevertheless, the model as such was not sufficient to understand the ice ages. To understand, completely, how the mechanism of ice age works, Milankovitch needed to know which seasons are crucial in triggering the glaciation process. The early theories were centered on very cold winters and accepted such severe winters as a decisive factor in accelerating glaciers (see Imbrie and Imbrie, 1979, p. 104). Milankovitch was skeptical about this hypothesis. However, being untrained in geological dynamics, he was far from abandoning this idea and putting an alternative approach. Yet, he was aware that he "had to solve" this "preliminary question of principal importance: which of the meteorological elements and which season was to be selected in the Ice-Period" (Milankovitch, 1941, p. 414).

In fact, the solution to this preliminary problem was proposed already by many scientists of the time. At the beginning of the 19<sup>th</sup> century, the German geologist and paleontologist Leopold von Buch (1774–1853) laid the foundation for this answer in Scandinavia, where he conducted his research on climatology. A main conclusion drawn from this research was the causal relationship between snow line levels and summer temperatures. This conclusion was in conflict with the common view that winters determine the snow line level. Nearly five decades later, Scottish physicist and glaciologist James David Forbes (1809–1868) made similar observations in Norway and supported Buch's claims with the following statement: "[observations support] the excellent generalization of von Buch, that it is the temperature of the summer months which determines the plane of perpetual snow" (Forbes 1853, p. 206). Shortly afterwards, Irish scientist Joseph John Murphy (1827–1894) maintained Forbes' line of thought by asserting "the idea that a long, cool summer and a short, mild winter are the most favorable conditions for glaciations" (Berger, 2021, p. 1729). As Murphy himself put, "we have plenty of observed data; and I think I can show that they all go to prove a cool summer to be what most promotes glaciation, while a cold winter has, usually, no effect on it whatever" (Murphy, 1869, p. 351). Nevertheless, it seems very likely that Milankovitch was unaware of these views and was in need of an urgent intellectual help.

Eventually, the required help came from the Russian–German geographer, and climatologist Wladimir Peter Köppen (1846–1940). He was aware of the works of Milankovitch and had read the monograph which is about the model of insolation. Soon after, he contacted Milankovitch. At the time, Köppen was researching on climate classification and climates of geological past with his son-in-law Wegener. Both researchers were experienced in the field, and they were familiar with almost all the relevant geological record as well as the seminal studies on ice ages. On a weekly basis, Köppen and



Milankovitch exchanged dozens of letters. Finally, the discussions ended up, when Köppen convinced him that the decisive factor on glaciation was milder summers, not colder winters as usually thought. That did seem reasonable to Milankovitch.

525 After an exhaustive discussion off all the possibilities, Köppen answered the question by indicating that it is the diminution of heat during the summer half-year which is the decisive factor in glaciation... I therefore used Köppen's advice and directed my attention to the periods of cold summers (Milankovitch, 1941, p. 414).

Truly, the suggestion to focus on milder summers was productive because the key factor in the development of glaciation was whether or not glaciers are preserved. At the high latitudes of the northern hemisphere, large amount of snow were accumulating and the glaciers were being formed even in completely different past climatic conditions. However, when the  
530 summers were hotter in any hemisphere, the glaciers formed in the winter were starting to melt slightly. Therefore, while milder or relatively colder summers were preserving glaciers and hence culminating glaciation; hotter summers were setting the stage for interglacial periods (see also Oerlemans, 1991).

Milankovitch had already constructed a model that makes it possible to calculate the insolation amount for a particular given latitude of the Earth. In this early version of the model, the strategy was to determine the difference between present and past  
535 values of insolation. The same strategy will also be followed in the revised model. However, the initial model was constructed on the basis of astronomical half-years. An astronomical year (or tropical year) is measured generally from either vernal equinox to the next one; or from the autumnal equinox to the next one. Following this measurement, the lengths of the seasons are considered as consisting of two parts. For instance, the northern hemisphere experiences astronomical summer half-year from 21 March (vernal equinox) to 23 September (autumnal equinox), and astronomical winter half-year  
540 from 23 September to 21 March. According to Milankovitch, however, the astronomical calendar was inappropriate to follow the secular march of insolation precisely because the length of each half-year in a particular year varies due to the variations of astronomical elements, especially due to the orbital eccentricity. Thus, this parameter should be modified correctly. In this manner, he divided "the year into two equally long seasons, one of which –the caloric summer– comprises all days during which the irradiation at the given latitude is stronger than on any day of the other half-year, i.e. the caloric  
545 winter" (Berger 2021, p. 1729).

As it is understood, astronomical half-years are not proper means to demonstrate variations in past climate. Caloric-half years were introduced merely for this reason. Nevertheless, dividing astronomical year into two caloric half-years is in a sense problematic. To deal with this problem, Milankovitch offered two solutions, one analytical and one geometrical. In the following, I will discuss the latter solution (see Fig.5) –those readers who are interested in analytical solution should refer to  
550 *Kanon* (1941, pp. 275-278).

In the graph, we have the representation of the mean radiation  $w$  of a surface element at any arbitrary latitude ( $\varphi$ ) as a function of time. Insolation is plotted along the vertical axis ( $w$ ), while time is plotted on the horizontal axis ( $t$ ). The points

on the curve PFGQHKSF' represents the course of radiation at the latitude ( $\phi$ ). The points I, II, III, IV and I' on the timeline, respectively denotes the time of the vernal equinox, the summer solstice, the autumnal equinox, the winter solstice and the subsequent equinox. The astronomical summer half-year ( $T_s$ ) is represented by the segment I–III, and the astronomical winter half-year ( $T_w$ ) is represented by the segment III–I' (see Milankovitch 1941, 271–294).

To understand this solution, it would be sufficient to notice the plotted points QH and SF' on the curve. These are the last irradiation intervals of astronomical summer half-year ( $T_s$ ) and astronomical winter half-year ( $T_w$ ), respectively. When we compare QH with SF', however, we see that the final irradiation interval of  $T_s$  (i.e. QH) is smaller than the final irradiation interval of  $T_w$  (i.e. SF'). Like in the astronomical calendar, each half-year's duration is 182 days, 14 hours and 54 minutes, but differently a caloric winter-half year ( $\frac{1}{2}T$  on the right-hand side) comprises all the days receiving relatively less irradiation than any of the day in summer-half year ( $\frac{1}{2}T$  on the left-hand side).

Using the latest version of his model, the version in which both caloric half-years and canonic units included, Milankovitch (with the suggestion of Köppen) calculated past insolation amounts for the specific latitudes, particularly 55<sup>th</sup>, 60<sup>th</sup> and 65<sup>th</sup> parallels of the northern hemisphere. He obtained the intended results over months and transformed these into fictitious latitudinal oscillations. The reason for that was to see a geometrical picture of the past summer insolation variations for the northern latitude of 65°. The created graph was an equivalent graph which showed the equivalent values of summer insolation amount for 65° N throughout the past 600.000 years (Fig. 6).

As seen in the graph, the past equivalent insolation values of 65° N are shown by curves for comparison. For example, about 115.000 years ago, the summer insolation amount of 65° N was almost equal to the today's summer insolation amount of 74° N. This could be evaluated as an indication of a relatively cold summer for the northern hemisphere. In the opposite cases of lower equivalent geographical latitude, the same latitude would of course have relatively warmer temperature averages in the summers.

Soon after completing his work, Milankovitch sent this radiation curve graph to Köppen. The graph must have been intriguing for Köppen, as it was in agreement with the findings of two German geographers Albrecht Penck (1858–1945) and Eduard Brückner (1862–1927). Penck and Brückner were researching on Alpine glaciers and about 15 years before Milankovitch's work, they identified four great glacial periods in Earth's history by examining successive gravels, plant remains and moraines in the European Alps (Anderson, Goudies and Parker, 2013 p. 8). They displayed these different glacial periods in a graph and named them chronologically as Günz, Mindel, Riss, and Würm in their 1909 book *Die Alpen im Eiszeitalter* (Fig. 7).

The close agreement of the two graphs was a sign of victory for Milankovitch. Moreover, with this new graph, Milankovitch provided a more precise graph compared to the old one. As seen in Fig. 6, the nine striped low points of the curve display colder summers which correspond to glacial periods occurred in the 589<sup>th</sup>, 548<sup>th</sup>, 475<sup>th</sup>, 434<sup>th</sup>, 231<sup>st</sup>, 187<sup>th</sup>, 116<sup>th</sup>, 72<sup>nd</sup>, and 22<sup>nd</sup> Millennia BP. Günz, Mindel and Riss, each includes two, and the last significant cold period Würm includes three low points. Clearly, Penck and Bruckner scheme does not have this preciseness.

## 7 Conclusion

On Milankovitch's part, the goal had been finally reached. His work "based upon exact science" passed into "the sphere of the descriptive natural sciences" and made the "link between celestial mechanics and geology" (Milankovitch, 1941, xvi). In short, exact science and descriptive science met by means of geological research on the one side and astronomical  
590 computation on the other. Furthermore, by constructing a mathematical model based on the orbital variations of Earth, it became possible to trace past climatic variations and also to predict the future glacial or interglacial periods. For example, the model predicts that, without human effects, interglacial period which we are still in may end about 50.000 years (see Berger and Loutre, 2002; Gajic 2019, p. 235, and also Paillard 2010).

Although, in 1924, Köppen and Wegener confidently published radiation curves in their book *Die Klimate der geologischen*  
595 *Vorzeit*, some remained sceptical about the validity of Milankovitch's model until the 1970's. In the following years, however, Hays *et al.* (1976) published an article on the close link between climatic changes and the Earth's orbital variations. Based on the oxygen isotope records provided by deep ocean sediment cores, they came to the conclusion that orbital changes induced climatic change in the past 500.000 years. In their terms, orbital cycles of the Earth should be understood as pacemakers of the ice ages. Eventually, this could be counted as a clinching victory for Milankovitch's model.  
600 In other words, Milankovitch's work was not merely providing a basis for further studies on mathematical climatology; it was also reliable in estimating the relationship between orbital variations and climatic changes, in a robust manner.

**Data availability.** No data sets were used in this article.

605 **Competing interests.** The author declares that he has no conflict of interest.

## Acknowledgments.

I would like to express my sincere gratitude to Kevin Hamilton, the handling topical editor, for his careful corrections and reasonable suggestions. Further, I would like to acknowledge the Anonymous Referee for her/his useful remarks and positive  
610 comments. I also remain grateful to Andreas Schmittner; his insightful and helpful comments did much to improve this article. Special thanks to Z. Bora Ön for his careful reading of the article and for making a number of constructive suggestions that improved substantially the initial manuscript. I wish to thank also to Jan Mangerud for kindly reminding me about recent works on astronomical theory of ice ages.

615 **References**

Adhémar, J. A.: Révolutions de la mer, Paris: Carilian–Goeury et V. Dalmont, 1842.

Agassiz, L.: On the Erratic Blocks of the Jura, The Edinburgh New Philosophical Journal XXIV (Oct. 1837–April 1838) pp.176–179, 1838.

Agassiz, L.: Early Discoverers III A Letter from Agassiz on Glaciation, Journal of Glaciology, 1(5), p. 294. doi:10.3189/002214349793702610, 1871/1949.

620 Agassiz, E. C.: Louis Agassiz. His Life and Correspondence, vol. I &II, Houghton. Boston: Mifflin & Company, 1886.

Andersen, B. G.: “Jens Esmark – a pioneer in glacial geology.” Boreas, 21(1), pp. 97–102. doi: <https://doi.org/10.1111/j.1502-3885.1992.tb00016.x>, 1992.

Anderson, D. E., Goudies, A. S. and Parker, A. G.: Global Environments through the Quaternary: Exploring Environmental Change, UK: Oxford University Press, 2013.

625 Bard, E.: Greenhouse effect and ice ages: historical perspective, C. R. Geoscience 336, 2004.

Barry, R. and Yew Gan, T.: The Global Cryosphere: Past, Present, and Future, Cambridge: Cambridge University Press, , 2011.

Berger, A.: A brief history of the astronomical theories of Paleoclimates, in: Berger, A., Mesinger, F., Sijacki, D. (eds), Climate Change: Inferences from Paleoclimate and Regional Aspects, Wien: Springer-Verlag, doi: 10.1007/978-3-7091-0973-1, 2012.

630 Berger, A.: Milankovitch, the father of paleoclimate modeling, Clim. Past (17), pp. 1727–1733, <https://doi.org/10.5194/cp-17-1727-2021>, 2021.

Berger, A. and Loutre, M. F.: An Exceptionally Long Interglacial Ahead?, Science 297(1287), pp.1287–1288, doi: 10.1126/science.1076120, 2002.

635 Bol'shakov, V., Kapitsa, A., & Rees, W. G.: James Croll: A scientist ahead of his time. Polar Record, 48(2), 201-205. doi:10.1017/S0032247411000301, 2012.

Boylan, P. J.: Lyell and the Dilemma of Quaternary glaciation, in: Blundell, D. J. and Scott, A. C. (eds) Lyell: the Past is the Key to the Present, London: Geological Society Special Publications, 143, 145-159, 2018.

640 Croll, J.: On the physical cause of the change of climate during geological epochs. The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science (28), 121-137. 1864.

Croll, J.: Climate And Time in Their Geological Relations: A Theory of Secular Changes of the Earth's Climate. London: Daldy, Isbister & Co., 1875.

Croll, J.: The Philosophical Basis of Evolution. London: E. Stanford., 1890.

645 Davis, O.: Ice Ages, in: Encyclopedia of Life Sciences eLS. John Wiley & Sons Ltd, doi: 10.1038/npg.els.0001635, 2001.

- Davies, G.: Early Discoverers XXVI: Another Forgotten Pioneer of the Glacial theory James Hutton (1726–97), *Journal of Glaciology*, 7(49), pp. 115-116. doi:10.3189/S0022143000020451, 1968.
- Eldredge, S. and Biek, B.: Ice ages –what are they and what causes them? *Survey Notes*, 42(3), Retrieved from [https://geology.utah.gov/wp-content/uploads/ice\\_ages1](https://geology.utah.gov/wp-content/uploads/ice_ages1), 2010.
- 650 Evans, E. P.: The Authorship of the Glacial Theory, *The North American Review*, vol. 145 (368), 1887.
- Forbes, J. D.: Norway and its glaciers, visited in 1851 - followed by journals of excursions in the high Alps of Dauphiné, Berne and Savoy, Edinburgh: A. and C. Black, 1853.
- Gajic, N.: The curious case of the Milankovitch calendar, *Hist. Geo Space. Sci.*, 10, 235–243, <https://doi.org/10.5194/hgss-10-235-2019>, 2019.
- 655 Grubic, A.: The astronomic theory of climatic changes of Milutin Milankovich, *Episodes*, vol. 29(3), 2006.
- Hay, W.W.: Tectonics and climate. *Geol Rundsch*, 85, 409–437, <https://doi.org/10.1007/BF02369000>, 1996.
- Hays, J. D., Imbrie, J. and Shackleton, N. J.: Variations in the Earth's Orbit: Pacemaker of the Ice Ages, *Science*, 194, 4270, pp. 1121-1132, 1976.
- Hestmark, G.: Jens Esmark's mountain glacier traverse 1823 – the key to his discovery of Ice Ages, *Boreas* 47(1), pp. 1-10, doi: <https://doi.org/10.1111/bor.12260>, 2018.
- 660 Imbrie J. and Imbrie K. P.: *Ice Ages: Solving the Mystery*, USA: Enslow Publishers, 1979.
- Janc, N., Gavrilov, M. B., Marković, S. B., Benišek, V. P., Popović, L. C. and Benišek, V.: Milutin Milanković and Associates in the Creation of the “Kanon”, in: Popović, L. Č., Srećković, V. A., Dimitrijević, M. S. and Kovačević, A. (eds), *Proceedings of the XII Serbian-Bulgarian Astronomical Conference "Publications of the Astronomical Society “Rudjer Bošković”*, No 20, pp. 123-128, 2020.
- 665 Langley, S. P.: The “Solar Constant” and Related Problems, *The Astrophysical Journal* (17), pp. 89–99, doi: [10.1086/140999](https://doi.org/10.1086/140999), 1903.
- Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records, *Paleoceanography*, 20, PA1003, doi: [10.1029/2004PA001071](https://doi.org/10.1029/2004PA001071), 2005.
- 670 Lutgens, F.K., Tarbuck, E. J. and Tasa, D.: *Essentials of Geology*, 11th Edition, USA: Prentice Hall, 2012.
- Maslin, M. A. and Ridgwell, A. J.: Mid-Pleistocene revolution and the eccentricity myth, in: Head, M. J. and Gibbard, E. L. (eds) *Early-Middle Pleistocene Transitions: The Land-Ocean Evidence*, London: Geological Society Special Publications, 247, 19-34, 2005.
- Milankovitch, M.: *Kanon der Erdbestrahlung und seine Anwendung auf das Eiszeitenproblem*, Belgrade: Royal Serbian Academy Special Publications, vol. 132, (Canon of insolation and the ice-age problem, English translation by Israel Program for Scientific Translations, Jerusalem, 1969), 1941.
- 675 Murphy, J. J.: On the nature and cause of the glacial climate, *Quarterly Journal of Geological Society* (25), pp. 350–356, 1869.
- Oerlemans, J.: The role of ice sheets in the Pleistocene climate, *Norsk Geologisk Tidsskrift*, vol. 71, pp. 155-161, 1991.

- 680 Paillard, D.: Climate and the orbital parameters of the Earth, *Comptes Rendus Geoscience*, 342 (4–5), pp. 273–285, doi: <https://doi.org/10.1016/j.crte.2009.12.006>, 2010.
- Paillard, D.: Quaternary glaciations: from observations to theories, *Quaternary Science Reviews* 107, doi: <https://doi.org/10.1016/j.crte.2004.02.005>, 2015.
- Petrovic, A.: Canon of Eccentricity: How Milankovic Built a General Mathematical Theory of Insolation”, in: Berger, A.,  
 685 Mesinger, F., Sijacki, D. (eds), *Climate Change: Inferences from Paleoclimate and Regional Aspects*, Wien: Springer-Verlag, doi: 10.1007/978-3-7091-0973-1\_9, 2012.
- Rohling, E. and PALAEOSSENS Project Members: Making sense of palaeoclimate sensitivity, *Nature* 491, 683–691, doi:10.1038/nature11574, 2012.
- Seylaz, L.: Early Discoverers XV: A Forgotten Pioneer of the Glacial Theory John Playfair (1748–1819). *Journal of*  
 690 *Glaciology*, 4(31), pp. 124–126. doi:10.3189/S002214300001827X, 1962.
- Thompson, R.: Croll, feedback mechanisms, climate change and the future, *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 112(3-4), 287–304. doi:10.1017/S1755691021000153, 2021.




Figure 1: The chart shows five major ice ages during the past 2.4 billion years. Adapted from Utah Geological Survey Notes, Eldredge S. and Biek B. (2010) [Adapted from Saltzman (2002)], © Copyright –Utah Geological Survey– State of Utah. May, 15, 2017.





Figure 2: A. The supercontinent Pangaea existed approximately between 350–260 million years ago. Ice sheets are located close to the south pole of the Earth. B. Present position of the continents. Old ice sheet evidences are located on different land masses. Reprinted from Essentials of Geology (p. 282) by Lutgens, F.K., Tarbuck, E. J. and Tasa, D., 2012, U.S.A.: Prentice Hall. Copyright ©2012, 2009, 2006 by Pearson Education, Inc.

710




715


**Figure 3** The figure shows all astronomical elements that have an influence in modulating the amount of insolation. Reprinted from *Kanon der Erdbestrahlung und seine Anwendung auf das Eiszeitenproblem* (p. 247) by Milankovitch, M., 1941, Belgrade: Royal Serbian Academy Special Publications.

720







**Figure 4: A.** The distances from the Sun at perihelion and aphelion for the Earth today. **B.** The maximum, minimum and present tilt angles for the Earth. **C.** The wobbling motion of the Earth.



**Figure 5: The annual march of the irradiation for any arbitrary latitude. Reprinted from *Kanon der Erdbestrahlung und seine Anwendung auf das Eiszeitenproblem* (p. 272) by Milankovitch, M., 1941, Belgrade: Royal Serbian Academy Special Publications.**



**Figure 6: The equivalent graph which shows the variations of the summer insolation for  $65^\circ N$ . Reprinted from *Kanon der Erdbestrahlung und seine Anwendung auf das Eiszeitenproblem* (p. 415) by Milankovitch, M., 1941, Belgrade: Royal Serbian Academy Special Publications.**



**Figure 7:** Penck and Brückner’s scheme of glacial/interglacial periods in the Alpines. Reprinted from *Kanon der Erdbestrahlung und seine Anwendung auf das Eiszeitenproblem* (p. 417) by Milankovitch, M., 1941, Belgrade: Royal Serbian Academy Special Publications.