

# 1 Atmospheric electricity observations by Reinhold Reiter 2 around Garmisch-Partenkirchen

3  
4 R. Giles Harrison<sup>a</sup> and Kristian Schlegel<sup>b</sup>  
5

6 <sup>a</sup>Department of Meteorology, Earley Gate, University of Reading, Reading, RG6 6ET UK

7 <sup>b</sup>Copernicus Gesellschaft, 3403 Göttingen, Germany  
8

9 Correspondence to: Giles Harrison ([r.g.harrison@reading.ac.uk](mailto:r.g.harrison@reading.ac.uk))

10  
11 submitted to *History of Geo and Space Sciences* (Atmospheric Electrical Observatories Special Issue)  
12  
13

14 **Abstract** Atmospheric electricity measurements were made at several sites close to Garmisch-Partenkirchen  
15 during four decades from 1950 to 1990 by Dr Reinhold Reiter, together with other environmental measurements.  
16 The quantities determined include the atmospheric potential gradient, the vertical current and the ion  
17 concentrations, and observations made at the Mount Wank site (1780 m, 47° 30' N, 11° 09' E) from 1st August  
18 1972 to 31st December 1983 are available in digital form.


19  
20 Keywords: Potential Gradient, conduction current; global circuit;  
21

## 22 1. Introduction

23 Motivated by his interest in the influence of atmospheric electric processes on humans,  
24 Reinhold Reiter (1920-1998) started atmospheric electricity measurements in the early 1950s.  
25 Past measurements of atmospheric electricity are increasingly studied internationally (Aplin,  
26 2020), because of widening interest in the global atmospheric electric circuit and its relevance  
27 to climate (e.g. Nicoll et al., 2019). Data obtained in clean air conditions are of particular  
28 importance, such as from mountain sites. The atmospheric electrical quantities obtained by  
29 Reiter within a sustained campaign of environmental measurements frequently fulfilled the  
30 clean air requirements.  
31

32 **Reiter** began with various measuring sites in Munich and southern Bavaria, probably to allow  
33 intercomparisons. Later, he concentrated on measurements undertaken at Garmisch-  
34 Partenkirchen, on the nearby Wank and Zugspitze mountains and onboard an instrumented  
35 passenger cable car moving regularly between the Eibsee and the Zugspitze summit. (The  
36 locations of these sites are shown in Fig. 1).

37



38

39 Figure 1: Area around Garmisch-Partenkirchen (southern Bavaria, Germany) **with scale** and  
40 the observational sites marked: Wank (square, upper right), Central Institute (triangle) and  
41 Zugspitze (star, lower left). The cable car runs almost directly north from the Zuspitze  
42 summit to the right hand shore of lake Eibsee (map adapted from Digitale Topographische  
43 Karte 1 : 100.000 (c) Bayerische Vermessungsverwaltung 2022, thanks to Martin Fasbender).

44

45 To undertake this, Reiter founded a privately-funded research institute, the *Physikalisch-*  
46 *bioklimatische Forschungsstelle in Garmisch-Partenkirchen* which was incorporated as the  
47 *Fraunhofer-Institut für Atmosphärische Umweltforschung (IFU)* in the Fraunhofer Society in  
48 1962. He led this institute as its director until his retirement in 1985. In 2002 this institute  
49 became part of the Institut für Meteorologie und Klimaforschung Atmosphärische  
50 Umweltforschung (IMK-IFU), and Campus Alpin of the Karlsruher Institut für Technologie  
51 (KIT).

52

53 Reinhold Reiter passed away on 24 September 1998 and a detailed memorial article was  
54 published by Weihe (1999). It is understood that some possessions were bequeathed to Ettal  
55 Abbey, a Benedictine monastery in Bavaria.

## 56 **2. Measurement locations**

57 Reiter's principal scientific motivations were to investigate biometeorological responses to  
58 atmospheric variables such as the concentrations of small ions, and to study short-term solar-  
59 terrestrial influences on the global circuit. This may be reflected in the choice of mountain sites  
60 for the measurements, which brought the possibility of low pollution conditions and least local  
61 disturbances.

62

63 The Garmisch-Partenkirchen measurements were obtained at permanent sites on the Zugspitze  
64 (2964 m altitude) and Wank (1780 m) mountains, and at an additional site known as the Central  
65 Research Institute, on the valley floor (740 m). A novel feature was the use of the cable car  
66 connecting the Zugspitze and a ground station close to lake Eibsee, instrumented to carry  
67 sensors in a regular path, sometimes passing repeatedly through fog and cloud layers. Vertical  
68 profiles of ozone were obtained using this approach (Reiter, 1991).

## 69 **3. Apparatus**

70 Customised instruments and systems were devised for the atmospheric electrical  
71 measurements. A primary quantity studied was the vertical potential gradient (PG). On Wank,  
72 as well as on the cable car, a radioactive collector probe was used, connected to a high  
73 impedance electrometer amplifier. The PG sensing probe was heated, and its physical  
74 construction refined during a long period of operation in mountain conditions, especially  
75 precipitation. The atmospheric conductivity was measured with an aspirated Gerdien  
76 condenser. A further measurement of the PG was made using an electrostatic field mill, and  
77 the air-earth current with a wire antenna. A special device was developed for measuring the  
78 space charge and, simultaneously, the natural radioactivity in the air. Beyond the usual fair  
79 weather measurements, the precipitation current density was obtained with an electric rain  
80 gauge. All these instruments and corresponding results are described in papers (Reiter, 1977a,  
81 b), and Reiter's textbook (Reiter, 1992).

## 82 **4. Data recovery**

83 Some of the measurements from the Bavarian Alps have previously been made available on a  
84 CDROM, which was originally distributed through the collaborative network provided by the  
85 SPECIAL scientific community (Rycroft and Füllekrug, 2004). These data values were

86 retrieved from magnetic tapes in summer 2000, with the help of one of Reiter's collaborators.  
 87 They provide hourly values from the Wank site (1780m, 47°30'N, 11°09'E), and span 1st  
 88 August 1972 to 31st December 1983. The wide range of quantities recorded is summarised in  
 89 Table 1, with the atmospheric electricity quantities identified.

90

91 **Table 1. Quantities recorded on Mount Wank (1972-1983)**

<i>Description of measured quantity</i>	<i>Symbol used in dataset</i>
<b>Meteorological and Environmental</b>	
air temperature	T
relative humidity	RF
water vapor partial pressure	E
specific humidity unit	SF
potential temperature	TH
equivalent potential temperature	THE
wind speed	WG
wind direction	WR
Sunshine duration	SD
Global solar irradiance	GS
Sky radiation	HS
UV intensity	UV
<b>Atmospheric Electrical</b>	
Electric field	F
Zero crossing of F	DU
Vertical current	I
Positive ion concentration	N+
Negative ion concentration	N-
Total ion concentration	SN
Positive ion conductivity	L+
Negative ion conductivity	L-
Total ion conductivity	SL


Number concentration of condensation nuclei	K1, K2, K3
---	------------

92

93 **5. Discussion**

94 The PG measurements obtained were over a sufficiently extended period to provide statistical  
 95 support for suspected solar effects on the lower atmosphere (Reiter, 1977b), which was a major  
 96 topic of research interest in the 1970s (e.g. Olson, 1971). **Due to these effects emerging, it is**  
 97 **likely that the local influences are sufficiently small that global atmospheric electric circuit**  
 98 **variations can also be retrieved.**

99




100

101 **Figure 2. Upper panel: Median hourly Potential Gradient (PG) across all months of the year**  
 102 **from Mount Wank, with the relative variation from Cruise VII of the Carnegie overplotted.**  
 103 **Lower panel: Hourly median PG by month from Mount Wank, using values for 1976-1983.**

104

105 Fig 2 provides a summary of the seasonal and diurnal variation in the PG at the site using data  
 106 from 1976 onwards, which is the longest period of consistent data following an unexplained  
 107 step change in the mean values. The upper panel of fig 2 shows the hourly variation across all  
 108 months, which is compared with the well-known global circuit “Carnegie curve” variation.  
 109 Although there are discrepancies in detail, perhaps arising from local meteorological factors or


110 uneven sampling, the (Pearson) correlation between hourly values of the Carnegie curve and  
 111 the Mount Wank PG is 0.96. The probability  $p$  that this is due to chance is small ( $p < 0.001$ ),  
 112 using the method of Ebisuzaki (1991) which accounts for serial correlation. The lower panel  
 113 of fig 2 shows the diurnal variations by month, in which the Carnegie curve is evident more  
 114 strongly in the second half of the year. Some values around midnight UTC are absent.



115  
 116 Figure 3. (a) Count of hourly samples of Mount Wank PG values 1976-1983 and (b)  
 117 distribution of all the hourly PG values obtained.

118  
 119 Fig 3 summarises the sampling and the distribution of values obtained. From fig 3a it can be  
 120 seen there are far fewer values for midnight and midday than for any of the other hours. It is  
 121 not clear why this is, but both midnight and midday occur first in each line of values in the data  
 122 files, so it might be a data processing artifact. A similar pattern of missing values is found for  
 123 some other measured quantities in the data files. Fig 3b presents the combined hourly PG data  
 124 as a histogram: the median is 84 Vm<sup>-1</sup>, and interquartile range 58 Vm<sup>-1</sup> to 119 Vm<sup>-1</sup>.

125  
 126 Fig 4 demonstrates the consistency evident between annual variations in PG measurements  
 127 from Mount Wank, and those made at Lerwick, Shetland (Harrison and Riddick, 2022), for  
 128 Decembers which have values available digitally. Some of the variations observed at Shetland  
 129 are thought to arise from the El Niño-Southern Oscillation (Harrison et al, 2022), in turn  
 130 modifying the global distribution of current-generating storms. Fig 4a shows the values as a  
 131 time series. Although there is a trend in the Mount Wank data (Harrison, 2004), fig 4b shows  
 132 the correlation between the two short series of values. This is consistent with the global circuit  
 133 providing the common variations occurring at both sites.



134

135 **Figure 4. Annual December mean PG values for Mount Wank compared with those for**  
 136 **Lerwick, following Harrison (2004), as (a) time series and (b) a scatterplot. The correlation**  
 137 **coefficient  $r$  in (b) is  $r = 0.74$  ( $p = 0.03$ ).**

138

139 **Combined with the Carnegie curve agreement of fig 2, fig 4 further supports the value of the**  
 140 **Mount Wank PG data for studying global circuit effects.**

## 141 **6. Conclusions**

142 Atmospheric electricity and other environmental measurements were made in the Bavarian  
 143 Alps over a long period, from which a series of hourly measurements for much of the 1970s is  
 144 available digitally. In the PG data **from the Mount Wank site**, the presence of global and solar-  
 145 terrestrial signals is apparent, which indicates the likely wider applicability of the  
 146 measurements. The endeavours at the Garmisch-Partenkirchen sites deserve to be more widely  
 147 known.

148

## 149 **Data availability**

150 The 1972-1983 Wank dataset is openly accessible through the University of Reading's  
 151 Research Data Archive, at <https://doi.org/10.17864/1947.000445> . **(The Lerwick December**  
 152 **data is available at <https://doi.org/10.17864/1947.000409> ).**

153

154

155 **Author Contributions**

156 The authors jointly drafted the manuscript.

157

158 **Competing interests**

159 Kristian Schlegel is an editorial board member of HGSS. There are no other competing  
160 interests.

161

162 **References**

163 Aplin, K. L.: Introduction to the special issue “Atmospheric electrical observatories”, *Hist.*  
164 *Geo Space. Sci.*, 11, 137–138, <https://doi.org/10.5194/hgss-11-137-2020> , 2020.

165 **Ebisuzaki, W.: A method to estimate the statistical significance of a correlation when the data**  
166 **are serially correlated. *Journal of Climate*, 10(9), 2147–2153. [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0442(1997)010<2147:AMTETS>2.0.CO;2)**  
167 **[0442\(1997\)010<2147:AMTETS>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<2147:AMTETS>2.0.CO;2), 1997.**

168 Harrison, R. G.: Long-range correlations in measurements of the global atmospheric electric  
169 circuit, *J. Atmos. Solar-Terrestrial Phys.*, 66, <https://doi.org/10.1016/j.jastp.2004.05.001> ,  
170 2004.

171 Harrison, R. G., Riddick J. C.: Atmospheric electricity observations at Lerwick Geophysical  
172 Observatory, *Hist. Geo Space. Sci.*, 13, 133–146, <https://doi.org/10.5194/hgss-13-133-2022> ,  
173 2022.

174 **Harrison, R.G., Nicoll, K.A., Joshi, M., Hawkins, E.: Empirical evidence for multidecadal**  
175 **scale Global Atmospheric Electric Circuit modulation by the El Niño-Southern Oscillation**  
176 ***Environ Res Lett* 17, 124048, <https://doi.org/10.1088/1748-9326/aca68c> 2022**

177 Nicoll, K. A., Harrison, R. G., Barta, V., Bor, J., Brugge, R., Chillingarian, A., Chum, J.,  
178 Georgoulas, A. K., Guha, A., Kourtidis, K., Kubicki, M., Mareev, E., Matthews, J.,  
179 Mkrtchyan, H., Odzimek, A., Raulin, J.-P., Robert, D., Silva, H. G., Tacza, J., Yair, Y., and  
180 Yaniv, R.: A global atmospheric electricity monitoring network for climate and geophysical  
181 research, *J. Atmos. Solar-Terrestrial Phys.*, 184, 18–29,  
182 <https://doi.org/10.1016/j.jastp.2019.01.003>, 2019.

183 Olson, D. E.: The evidence for auroral effects on atmospheric electricity, *Pure Appl.*  
184 *Geophys. PAGEOPH*, 84, 118–138, <https://doi.org/10.1007/BF00875461>, 1971.

185 Reiter, R.: Review of the History, Activities and Basic Facilities of the Institute for  
186 Atmospheric Environmental Research, in: *Electrical Processes in Atmospheres*, 759–803,  
187 1977a.

188 Reiter, R.: The electric potential of the ionosphere as controlled by the solar magnetic sector  
189 structure. Result of a study over the period of a solar cycle, *J. Atmos. Terr. Phys.*, 39,  
190 [https://doi.org/10.1016/0021-9169\(77\)90048-4](https://doi.org/10.1016/0021-9169(77)90048-4), 1977b.

191 Reiter, R.: On the mean daily and seasonal variations of the vertical ozone profiles in the



- 192 lower troposphere. Atmos Environ, 25A, 9, pp1751-1757, 1991.
- 193 Reiter, R.: Phenomena in atmospheric and environmental electricity, Elsevier Science  
194 Publishers, Amsterdam (572pp), 1992.
- 195 Rycroft, M. J. and Füllekrug, M.: The initiation and evolution of SPECIAL, J. Atmos. Solar-  
196 Terrestrial Phys., 66, <https://doi.org/10.1016/j.jastp.2004.05.013>, 2004.
- 197 Weihe, W. H.: In memoriam Reinhold Reiter 17 November 1920-24 September 1998, Int. J.  
198 Biometeorol., 43, <https://doi.org/10.1007/s004840050122>, 1999.